arXiv:0809.3206v1 [hep-th] 18 Sep 2008

Basic properties of Fedosov and Riemannian supermanifolds

M. Asorey[a], B. Geyer[b], P.M. Lavrov[a,c] and O.V. Radchenko[c]

[a] *Departamento de Física Teórica, Facultad de Ciencias Universidad de Zaragoza,*
50009 Zaragoza, Spain
[b] *Institute of Theoretical Physics,Leipzig University,*
D-04109 Leipzig, Germany
[c] *Department of Mathematical Analysis,Tomsk State Pedagogical University,*
634041 Tomsk, Russia

We discuss some differences in the properties of both even and odd Fedosov and Riemannian supermanifolds.

A *Fedosov supermanifold* (M, ω, Γ) is defined as a symplectic supermanifold (M, ω) equipped with a symmetric connection Γ (or covariant derivative ∇) compatible with a given symplectic structure ω: $\omega\nabla = 0$.[1] In local coordinates $\{x^i\}$, $\epsilon(x^i) = \epsilon_i$, on the supermanifold M the symplectic structure is $\omega = \omega_{ij} dx^j \wedge dx^i$, $\quad \omega_{ij} = -(-1)^{\epsilon_i\epsilon_j}\omega_{ji}$ (Grassmann parity of the symplectic structure, $\epsilon(\omega)$, is equal to 0 for even structure and 1 for odd structure) and the compatibility condition is $\omega_{ij}\nabla_k = \omega_{ij,k} - \Gamma_{ijk} + \Gamma_{jik}(-1)^{\epsilon_i\epsilon_j} = 0$ where $\Gamma_{ijk} = \omega_{in}\Gamma^n_{\ jk}$, $\quad \epsilon(\Gamma_{ijk}) = \epsilon(\omega) + \epsilon_i + \epsilon_j + \epsilon_k$ and $\Gamma^i_{\ jk}$ are components of the connection Γ. Notice that for a given symplectic structure ω there exists a large family of connections satisfying the compatibility condition.

The curvature tensor field $R^i_{\ mjk}$ is defined in terms of the commutator of covariant derivatives, $[\nabla_i, \nabla_j] = \nabla_i\nabla_j - (-1)^{\epsilon_i\epsilon_j}\nabla_j\nabla_i$, whose action on a vector field T^i is $T^i[\nabla_j, \nabla_k] = -(-1)^{\epsilon_m(\epsilon_i+1)} T^m R^i_{\ mjk}$.

It is convenient to describe the basic properties of Fedosov supermanifolds in terms of the symplectic curvature tensor $R_{ijkl} = \omega_{in}R^n_{\ jkl}$, $\quad \epsilon(R_{ijkl}) = \epsilon(\omega) + \epsilon_i + \epsilon_j + \epsilon_k + \epsilon_l,$. This tensor obeys the symmetry properties

$$R_{ijkl} = -(-1)^{\epsilon_k\epsilon_l} R_{ijlk}\,, \quad R_{ijkl} = (-1)^{\epsilon_i\epsilon_j} R_{jikl}\,, \tag{1}$$

and satisfies the Jacobi identity

$$(-1)^{\epsilon_j\epsilon_l} R_{ijkl} + (-1)^{\epsilon_l\epsilon_k} R_{iljk} + (-1)^{\epsilon_k\epsilon_j} R_{iklj} = 0\,, \tag{2}$$

the Bianchi identity

$$(-1)^{\epsilon_k\epsilon_m} R_{ijkl;m} + (-1)^{\epsilon_l\epsilon_m} R_{ijmk;l} + (-1)^{\epsilon_k\epsilon_l} R_{ijlm;k} = 0\,, \tag{3}$$

---

[1] We use conventions and definitions adopted in [1, 2, 3].

and the special symplectic identity

$$R_{ijkl}+(-1)^{\epsilon_l(\epsilon_i+\epsilon_k+\epsilon_j)}R_{lijk}+(-1)^{(\epsilon_k+\epsilon_l)(\epsilon_i+\epsilon_j)}R_{klij}+(-1)^{\epsilon_i(\epsilon_j+\epsilon_l+\epsilon_k)}R_{jkli}=0. \qquad (4)$$

We see that there are no formal differences in the properties of even and odd Fedosov supermanifolds on the level of symplectic curvature tensor. With the curvature tensor, R_{ijkl}, and the inverse tensor field ω^{ij} ($\omega^{ij}=-(-1)^{\epsilon(\omega)+\epsilon_i\epsilon_j}\omega^{ji}$) of the symplectic structure ω_{ij}, one can construct the only tensor field of type $(0,2)$,

$$K_{ij}=\omega^{kn}R_{nikj}(-1)^{\epsilon_i\epsilon_k+(\epsilon(\omega)+1)(\epsilon_k+\epsilon_n)} \;=\; R^{k}{}_{ikj}\,(-1)^{\epsilon_k(\epsilon_i+1)}, \quad \epsilon(K_{ij})=\epsilon_i+\epsilon_j. \qquad (5)$$

This tensor satisfies the relations [1]

$$[1+(-1)^{\epsilon(\omega)}](K_{ij}-(-1)^{\epsilon_i\epsilon_j}K_{ji})=0\,, \qquad (6)$$

and is called the Ricci tensor. In the even case this tensor is symmetric whereas in the odd case there are not restrictions on its (generalized) symmetry properties. The scalar curvature tensor K is defined by the formula $K=\omega^{ji}K_{ij}(-1)^{\epsilon_i+\epsilon_j}$. From the symmetry properties of R_{ijkl}, it follows that

$$[1+(-1)^{\epsilon(\omega)}]K=0. \qquad (7)$$

Therefore as in the case of Fedosov manifolds [4], even Fedosov supermanifolds have vanishing scalar curvature K. However, for odd Fedosov supermanifolds this curvature is, in general, not vanishing. This fact was quite recently used in Ref. [5] to generalize the BV formalism [6].

A *Riemannian supermanifold* (M,g,Γ) is defined as a metric supermanifold (M,g) equipped with a symmetric connection Γ (or covariant derivative ∇) compatible with a given metric structure g: $g\nabla=0$. In local coordinates on the supermanifold M the metric structure is $g=g_{ij}dx^j dx^i, \quad g_{ij}=(-1)^{\epsilon_i\epsilon_j}g_{ji}$ (Grassmann parity of the metric structure, $\epsilon(g)$, is equal to 0 for even structure and 1 for odd structure) and the compatibility condition is $g_{ij}\nabla_k=g_{ij,k}-\Gamma_{ijk}-\Gamma_{jik}(-1)^{\epsilon_i\epsilon_j}=0$ where $\Gamma_{ijk}=g_{in}\Gamma^{n}{}_{jk}, \quad \epsilon(\Gamma_{ijk})=\epsilon(g)+\epsilon_i+\epsilon_j+\epsilon_k$ and $\Gamma^{i}{}_{jk}$ are components of the connection Γ. Notice that for a given metric structure g there exists the unique symmetric connection $\Gamma^{i}{}_{jk}$ which is compatible with a given metric structure,

$$\Gamma^{l}{}_{ki}=\frac{1}{2}g^{lj}\Big(g_{ij,k}(-1)^{\epsilon_k\epsilon_i}+g_{jk,i}(-1)^{\epsilon_i\epsilon_j}-g_{ki,j}(-1)^{\epsilon_k\epsilon_j}\Big)(-1)^{\epsilon_j\epsilon_i+\epsilon_j+\epsilon(g)(\epsilon_j+\epsilon_l)}, \qquad (8)$$

where g^{ij} is the inverse tensor field of the metric g_{ij} ($g^{ij}=(-1)^{\epsilon(g)+\epsilon_i\epsilon_j}g^{ji}, \quad \epsilon(g^{ij})=\epsilon(g)+\epsilon_i+\epsilon_j$).

The curvature tensor $\mathcal{R}_{ijkl}=g_{in}\mathcal{R}^{n}{}_{jkl}$ ($\epsilon(\mathcal{R}_{ijkl})=\epsilon(g)+\epsilon_i+\epsilon_j+\epsilon_k+\epsilon_l$), obeys the symmetry properties

$$\mathcal{R}_{ijkl}=-(-1)^{\epsilon_k\epsilon_l}\mathcal{R}_{ijlk}, \quad \mathcal{R}_{ijkl}=-(-1)^{\epsilon_i\epsilon_j}\mathcal{R}_{jikl}, \quad \mathcal{R}_{ijkl}=\mathcal{R}_{klij}(-1)^{(\epsilon_i+\epsilon_j)(\epsilon_k+\epsilon_l)} \qquad (9)$$

and satisfies the Jacobi identity (2) and the Bianchi identity (3). Again we find that on the level of the curvature tensor there are no differences in the basic properties of even and odd Riemannian supermanifolds.

From the curvature tensor $\mathcal{R}_{ijkl}$ and the inverse tensor field g^{ij} of the metric g_{ij} one can define the only tensor field of type $(0, 2)$:

$$\mathcal{R}_{ij} = \mathcal{R}^{k}{}_{ikj}(-1)^{\epsilon_k(\epsilon_i+1)} = g^{kn}\mathcal{R}_{nikj}(-1)^{(\epsilon_k+\epsilon_n)(\epsilon(g)+1)+\epsilon_i\epsilon_k}, \quad \epsilon(\mathcal{R}_{ij}) = \epsilon_i + \epsilon_j. \tag{10}$$

It is the generalized Ricci tensor with the following symmetry properties

$$R_{ij} = (-1)^{\epsilon(g)+\epsilon_i\epsilon_j} R_{ji} \tag{11}$$

depending on Riemannian supermanifolds to be even or odd. A further contraction defines the scalar curvature

$$\mathcal{R} = g^{ji}\mathcal{R}_{ij}\ (-1)^{\epsilon_i+\epsilon_j}, \quad \epsilon(\mathcal{R}) = \epsilon(g) \tag{12}$$

which, in general, is not equal to zero. Notice that for an odd metric structure the scalar curvature tensor squared is identically equal to zero, $\mathcal{R}^2 = 0$.

From the Bianchi identity one can deduce the following relation between the scalar curvature and the Ricci tensor

$$\mathcal{R}_{,i} = [1 + (-1)^{\epsilon(g)}]\mathcal{R}^{j}{}_{i;j}(-1)^{\epsilon_j(\epsilon_i+1)}, \tag{13}$$

which in the even case is nothing but $\mathcal{R}_{,i} = 2\mathcal{R}^{j}{}_{i;j}(-1)^{\epsilon_j(\epsilon_i+1)}$, i.e. the supersymmetric generalization of known relation in Riemannian geometry [7]. In the odd case $\mathcal{R}_{,i} = 0$. Therefore odd Riemann supermanifolds have constant scalar curvature, $\mathcal{R} = const.$

References

[1] B. Geyer and P.M. Lavrov, *Int. J. Mod. Phys.*, **A19**, 3195 (2004).

[2] P.M. Lavrov and O.V. Radchenko, *Theor. Math. Phys.*, **149**, 1474 (2006).

[3] M. Asorey and P.M. Lavrov, Fedosov and Riemannian supermanifolds, arXiv:0803.1591.

[4] I. Gelfand, V. Retakh and M. Shubin, *Adv. Math.*, **136**, 104 (1998); [dg-ga/9707024].

[5] I.A. Batalin and K. Bering, Odd Scalar Curvature in Field-Antifield Formalism, arXiv:0708.0400; Odd Scalar Curvature in Anti-Poisson Geometry, arXiv:0712.3699.

[6] I.A. Batalin and G.A. Vilkovisky, *Phys. Lett.*, **B102**, 27 (1981); *Phys. Rev.*, **D28**, 2567 (1983).

[7] L.P. Eisenhart, *Riemannian Geometry*, Princeton University Press, 1949.